

FD-RBF for Partial Integro-Differential Equations with a Weakly Singular Kernel

Jafar Biazar^{*}, Mohammad Ali Asadi

Department of Applied Mathematics, Faculty of Mathematical Science, University of Guilan, Rasht, Iran

Email address:

biazar@guilan.ac.ir (J. Biazar), eng.asadi@gmail.com (M. A. Asadi)

To cite this article:

Jafar Biazar, Mohammad Ali Asadi. FD-RBF for Partial Integro-Differential Equations with a Weakly Singular Kernel. *Applied and Computational Mathematics*. Vol. 4, No. 6, 2015, pp. 445-451. doi: 10.11648/j.acm.20150406.17

Abstract: Finite Difference Method and Radial Basis Functions are applied to solve partial integro-differential equations with a weakly singular kernel. The product trapezoidal method is used to compute singular integrals that appear in the discretization process. Different RBFs are implemented and satisfactory results are shown the ability and the usefulness of the proposed method.

Keywords: Partial Integro-Differential Equations (PIDE), Weakly Singular Kernel, Radial Basis Functions (RBF), Finite Difference Method (FDM), Product Trapezoidal Method

1. Introduction

Mathematical modeling of some scientific and engineering problems lead to partial integro-differential equations (PIDEs). In this paper, the following PIDE, with a weakly singular kernel is considered.

$$u_t(x, t) = \mu u_{xx}(x, t) + \int_0^t (t-s)^{-\frac{1}{2}} u_{xx}(x, s) ds, \quad 0 < x < 1, 0 \leq t \leq T \quad (1)$$

where $\mu \geq 0$, and is subject to the following boundary and initial conditions:

$$u(0, t) = u(1, t) = 0, \quad 0 \leq t \leq T,$$

$$u(x, 0) = g(x), \quad 0 \leq x \leq 1.$$

This type of integro-differential equations appears in some phenomena such as heat conduction in materials with memory [12, 22], population dynamics, and viscoelasticity [25, 6]. The numerical solution of PIDEs is considered by many authors [1, 2, 19, 20, 23, 27, 30].

PIDEs with weakly singular kernels have been studied in some papers. Numerical solution of a parabolic integro-differential equation with a weakly singular kernel by means of the Galerkin finite element method is discussed in [5]. A finite difference scheme and a compact difference scheme are presented for PIDEs, with a weakly singular kernel, in [28] and [21], respectively. A spectral collocation method is considered in [17] for weakly singular PIDEs. Also Quintic B-spline collocation method [31] and Crank-Nicolson/quasi-wavelets method [32]

are used for solving fourth order partial integro-differential equation with a weakly singular kernel and some others [10, 14].

In recent years, meshless methods, as a class of numerical methods, are used for solving functional equations. Meshless methods just use a scattered set of collocation points, regardless any relationship between the collocation points. This property is the main advantage of these techniques in comparison with the mesh dependent methods, such as finite difference and finite element. Since 1990, radial basis function methods (RBF) [13] are used as a well-known family of meshless methods to approximate the solutions of various types of linear and nonlinear functional equations, such as Partial Differential Equations (PDEs), Ordinary Differential Equations (ODEs), Integral Equations (IEs), and Integro-Differential Equations (IDEs) [7, 8, 11, 13, 16, 18, 24]. In the present work, for the first time derivatives, we use the Finite Difference (FD) scheme to discretize the equation which it makes a system of partial integro-differential equations. Then we use radial basis functions (RBFs) to solve this system. Recently FD-RBF method is used to solve some problems like nonlinear parabolic-type Volterra partial integro-differential equations [1], fractional-diffusion inverse heat conduction problems [33], and wave equation with an integral condition [34].

In this paper, FD-RBF methods are applied for numerical solution of PIDEs with a weakly singular kernel. Singular integrals, which appear in the method, are computed by the product trapezoidal integration rule.

The paper is organized as follows. In Section 2, the RBFs are introduced. Section 3, as the main part, is devoted to

solving weakly singular PIDEs, by finite difference and RBFs. An illustrative example is included in Section 4. A conclusion is presented in Section 5.

2. Radial Basis Functions

Interpolation of a function $u : \mathbb{R}^d \rightarrow \mathbb{R}$ by RBF can be presented as the following [4]

$$s_N(\mathbf{x}) := \sum_{i=0}^N \lambda_i \phi(\|\mathbf{x} - \mathbf{x}_i\|), \quad \mathbf{x} \in \mathbb{R}^d. \quad (2)$$

Where $\phi : [0, \infty) \rightarrow \mathbb{R}$ is a fixed univariate function, the coefficients $(\lambda_i)_{i=0}^N$ are real numbers, $(\mathbf{x}_i)_{i=0}^N$ is a set of interpolation points in \mathbb{R}^d , and $\|\cdot\|$ is the Euclidean norm.

Eq. (2) can be written as follows

$$s_N(\mathbf{x}) := \sum_{i=0}^N \lambda_i \phi_i(\mathbf{x}) = \Phi^T(\mathbf{x})\Lambda, \quad (3)$$

where

$$\begin{aligned} \phi_i(\mathbf{x}) &= \phi(\|\mathbf{x} - \mathbf{x}_i\|), \\ \Phi(\mathbf{x}) &= [\phi_0(\mathbf{x}), \phi_1(\mathbf{x}), \dots, \phi_N(\mathbf{x})]^T, \\ \Lambda &= [\lambda_0, \lambda_1, \dots, \lambda_N]^T. \end{aligned}$$

Consider $N+1$ distinct support points $(\mathbf{x}_j, u(\mathbf{x}_j))$, $j = 0, 1, \dots, N$. One can use interpolation conditions to find λ_i s by solving the following linear system

$$\mathbf{A}\Lambda = \mathbf{u},$$

in which

$$\mathbf{A} = [\phi(\|\mathbf{x}_j - \mathbf{x}_i\|)]_{i,j=0}^N,$$

and

$$\mathbf{u} = [u(\mathbf{x}_0), u(\mathbf{x}_1), \dots, u(\mathbf{x}_N)]^T$$

Table 1. Some well-known RBFs.

Name of the RBF	Definition
Gaussian	$\phi(r) = e^{-(cr)^2}$
Inverse Quadric	$\phi(r) = \frac{1}{r^2 + c^2}$
Hardy Multiquadric	$\phi(r) = \sqrt{r^2 + c^2}$
Inverse Multiquadric	$\phi(r) = \frac{1}{\sqrt{r^2 + c^2}}$
Cubic	$\phi(r) = r^3$
Thin Plate Spline	$\phi(r) = r^2 \log(r)$
Hyperbolic Secant	$\phi(r) = \text{sech}(\frac{r}{c})$

Some well-known RBFs are listed in Table 1, where the Euclidean distance r is real and non-negative, and c is a positive scalar, called the shape parameter.

Also the generalized Thin Plate Splines (TPS) are defined as the following:

$$\phi(r) = r^{2m} \log(r), \quad m = 1, 2, \dots$$

Some of RBFs are unconditionally positive definite (e.g. Gaussian or Inverse Multiquadrics) to guarantee that the resulting system is solvable, and some of them are conditionally positive definite. Although, some of RBFs are conditionally positive definite functions, polynomials are augmented to Eq. (2) to guarantee that the outcome interpolation matrix is invertible. Such an approximation can be expressed as follows

$$s(\mathbf{x}) = \sum_{i=0}^N \lambda_i \phi(\|\mathbf{x} - \mathbf{x}_i\|) + \sum_{i=1}^l \lambda_{N+i} p_i(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^d. \quad (4)$$

where $p_i(x)$, $i = 1, \dots, l$, are polynomials on \mathbb{R}^d of degree at most $m-1$, and $l = \binom{m-1}{d}$. Here l is the dimension of

the linear space Π_{m-1}^d of polynomials of total degree less than or equal to m with d variables.

Collocation method is used to determinate the coefficients $(\lambda_0, \lambda_1, \dots, \lambda_N)$ and $(\lambda_{N+1}, \lambda_{N+2}, \dots, \lambda_{N+l})$. This will produce $N+1$ equations at $N+1$ points. l additional equations is usually written in the following form

$$\sum_{i=0}^N \lambda_i p_j(\mathbf{x}_i) = 0, \quad j = 1, \dots, l \quad (5)$$

3. Application of FD-RBF Method

In this section we explain the process of solving PIDEs, with a weakly singular kernel, in the following form

$$u_t(x, t) = \mu u_{xx}(x, t) + \int_0^t (t-s)^{-\frac{1}{2}} u_{xx}(x, s) ds, \quad 0 < x < 1, \quad 0 \leq t \leq T, \quad (6)$$

where $u_t = \frac{\partial u}{\partial t}$, $u_{xx} = \frac{\partial^2 u}{\partial x^2}$, $\mu \geq 0$, with the following boundary and initial conditions

$$u(0, t) = u(1, t) = 0, \quad 0 \leq t \leq T, \quad (7)$$

$$u(x, 0) = g(x), \quad 0 \leq x \leq 1. \quad (8)$$

At first we introduce grid points $t_i = a + ih$, $i = 0, 1, \dots, M$, where $M \geq 1$ is an integer, and $h = (T - 0) / M$.

Considering (6) at point (x, t_i) , we have

$$u_t(x, t_i) = \mu u_{xx}(x, t_i) + \int_0^{t_i} (t_i - s)^{-\frac{1}{2}} u_{xx}(x, s) ds, \quad i = 1, \dots, M \quad (9)$$

As the finite difference technique, we have

$$u_i(x, t_i) \approx \frac{u(x, t_{i+1}) - u(x, t_{i-1})}{2h}.$$

Discretizing (9) by the θ -weighted method leads to

$$\frac{u(x, t_{i+1}) - u(x, t_{i-1})}{2h} = \mu(\theta u_{xx}(x, t_{i+1}) + (1-\theta)u_{xx}(x, t_i)) + \int_0^{t_i} (t_i - s)^{-\frac{1}{2}} (\theta u_{xx}(x, s+h) + (1-\theta)u_{xx}(x, s)) ds,$$

$i = 1, \dots, m-1$, and $\theta \in [0, 1]$. By using the notation $u^i(x) = u(x, t_i)$ we have

$$u^{i+1}(x) = u^{i-1}(x) + 2h\mu(\theta u_{xx}^{i+1}(x) + (1-\theta)u_{xx}^i(x)) + 2h \int_0^{t_i} (t_i - s)^{-\frac{1}{2}} (\theta u_{xx}(x, s+h) + (1-\theta)u_{xx}(x, s)) ds. \quad (10)$$

We now use the product trapezoidal integration technique, well addressed in [9], to approximate the integrals

$$\int_0^{t_i} (t_i - s)^{-\frac{1}{2}} u_{xx}(x, s+h) ds \approx \sum_{j=0}^i w_{ij} u_{xx}^{j+1}(x), \quad (11)$$

and

$$\int_0^{t_i} (t_i - s)^{-\frac{1}{2}} u_{xx}(x, s) ds \approx \sum_{j=0}^i w_{ij} u_{xx}^j(x), \quad (12)$$

where

$$w_{i0} = \frac{1}{h} \int_{t_0}^{t_i} \frac{t_i - s}{\sqrt{t_i - s}} ds, \\ w_{ij} = \frac{1}{h} \left(\int_{t_{j-1}}^{t_j} \frac{s - t_{j-1}}{\sqrt{t_i - s}} ds + \int_{t_j}^{t_{j+1}} \frac{t_{j+1} - s}{\sqrt{t_i - s}} ds \right), \quad j=1, 2, \dots, i-1 \\ w_{ii} = \frac{1}{h} \int_{t_{i-1}}^{t_i} \frac{s - t_{i-1}}{\sqrt{t_i - s}} ds.$$

Substituting (11) and (12) into (10), results in

$$u^{i+1}(x) = u^{i-1}(x) + 2h\mu(\theta u_{xx}^{i+1}(x) + (1-\theta)u_{xx}^i(x)) + 2h\theta \sum_{j=0}^i w_{ij} u_{xx}^{j+1}(x) + 2h(1-\theta) \sum_{j=0}^i w_{ij} u_{xx}^j(x),$$

or

$$u^{i+1}(x) - 2h\theta(\mu + w_{ii})u_{xx}^{i+1} = u^{i-1}(x) + 2h\mu(1-\theta)u_{xx}^i(x) + 2h(1-\theta)w_{i0}u_{xx}^0(x) + 2h \sum_{j=1}^i (\theta w_{i,j-1} + (1-\theta)w_{ij})u_{xx}^j(x), \quad (13)$$

for $i=1, \dots, m-1$. Let's approximate the function $u^i(x)$ in terms of RBFs, as follows

$$u^i(x) = \sum_{n=0}^N \lambda_{i,n} \phi_n(x) + \lambda_{i,N+1}x + \lambda_{i,N+2} = \Phi^T(x) \Lambda^i, \quad (14)$$

where

$$\phi_n(x) = \phi(\|x - x_n\|),$$

$x_n = a + nk$, $n = 0, 1, \dots, N$, are center points, $k = (b-a)/N$ and $N \geq 1$ is an integer,

$$\Phi(x) = [\phi_0(x), \dots, \phi_N(x), x, 1]^T \quad (15)$$

and $\Lambda^i = [\lambda_{i,0} \dots \lambda_{i,N} \lambda_{i,N+1} \lambda_{i,N+2}]^T$ is an unknown vector.

Collocating Eq. (14) at $(x_j)_{j=0}^N$ leads to the following system

$$u_j^i := u^i(x_j) = \sum_{n=0}^N \lambda_{i,n} \phi_n(x_j) + \lambda_{i,N+1}x_j + \lambda_{i,N+2}, \quad j=0, \dots, N$$

Two additional conditions, as mentioned in section 2, can be written as

$$\sum_{j=0}^N \lambda_{i,j} = 0, \quad (1)$$

$$\sum_{j=0}^N \lambda_{i,j} x_j = 0. \quad (2)$$

Eqs. (14), (16), and (17) can be written in the following matrix form

$$u^i = \Lambda \Lambda^i,$$

where

$$u^i = [u_0^i, \dots, u_N^i, 0, 0]^T, \quad \Lambda^i = [\lambda_{i,0} \dots \lambda_{i,N} \lambda_{i,N+1} \lambda_{i,N+2}]^T, \text{ and}$$

$$A = \begin{bmatrix} \phi_0(x_0) & \phi_1(x_0) & \dots & \phi_N(x_0) & x_0 & 1 \\ \phi_0(x_1) & \phi_1(x_1) & \dots & \phi_N(x_1) & x_1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \phi_0(x_N) & \phi_1(x_N) & \dots & \phi_N(x_N) & x_N & 1 \\ x_0 & x_1 & \dots & x_N & 0 & 0 \\ 1 & 1 & \dots & 1 & 0 & 0 \end{bmatrix} \quad (18)$$

By two times differentiation from Eq. (14), with respect to x , we obtain

$$u_{xx}^i = \sum_{n=0}^N \lambda_{i,n} \phi_n''(x) = \Psi^T(x) \Lambda^i, \quad (19)$$

Where

$$\Psi(x) = [\phi_0''(x), \phi_1''(x), \dots, \phi_N''(x), 0, 0]^T$$

Substituting Eqs. (14) and (19) in Eq. (13), leads to

$$\begin{aligned} & \Phi^T(x) \Lambda^{i+1} - 2h\theta(\mu + w_{ij}) \Psi^T(x) \Lambda^{i+1} \\ &= \Phi^T(x) \Lambda^{i+1} + 2h\mu(1-\theta) \Psi^T(x) \Lambda^i + 2h(1-\theta)w_{i0} \Psi^T(x) \Lambda^0 \\ &+ 2h \sum_{j=1}^i (\theta w_{i,j-1} + (1-\theta)w_{ij}) \Psi^T(x) \Lambda^j, \end{aligned} \quad (3)$$

for $i = 1, \dots, M-1$. So, we consider collocation points, x_n , $n = 0, 1, \dots, N$, to obtain the entries of the vectors of the coefficients Λ^i , $i = 1, \dots, M$ in Eq. (20). This leads to

$$\begin{aligned} & \mathbf{A} \Lambda^{i+1} - 2h\theta(\mu + w_{ij}) \mathbf{T} \Lambda^{i+1} = \mathbf{A} \Lambda^{i+1} + 2h\mu(1-\theta) \mathbf{T} \Lambda^i \\ &+ 2h(1-\theta)w_{i0} \mathbf{T} \Lambda^0 + 2h \sum_{j=1}^i (\theta w_{i,j-1} + (1-\theta)w_{ij}) \mathbf{T} \Lambda^j, \end{aligned} \quad (21)$$

where $i = 1, \dots, M-1$, and

$$T = \begin{bmatrix} 0 & \dots & 0 & 0 & 0 \\ \phi''_0(x_1) & \dots & \phi''_N(x_1) & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ \phi''_0(x_{N-1}) & \dots & \phi''_N(x_{N-1}) & 0 & 0 \\ 0 & \dots & 0 & 0 & 0 \\ 0 & \dots & 0 & 0 & 0 \\ 0 & \dots & 0 & 0 & 0 \end{bmatrix}$$

As the first step we must determine Λ^0 and Λ^1 .

Obviously, Λ^0 can be obtained by the initial condition

$$\begin{aligned} & u^0(x) = u(x, 0) = g(x) \\ \Rightarrow \mathbf{A} \Lambda^0 = \mathbf{u}_0 &= [g(x_0) \quad g(x_1) \quad \dots \quad g(x_N) \quad 0 \quad 0]^T. \end{aligned}$$

To approximate Λ^1 , we use

$$u_t(x, t_1) \approx \frac{u(x, t_1) - u(x, t_0)}{h}. \quad (22)$$

Substituting (22) into (9), similarly leads to

$$(\mathbf{A} - v\mathbf{T} - h\mu\mathbf{T}) \Lambda^1 = (\mathbf{A} + w\mathbf{T}) \Lambda^0, \quad (23)$$

where

$$v = \int_0^{t_1} \frac{s - t_1}{\sqrt{t_2 - s}} ds,$$

and

$$w = \int_0^{t_1} \frac{t_2 - s}{\sqrt{t_2 - s}} ds.$$

The convergence of RBF interpolation has been addressed by Buhmann [3, 4], and other researchers [15, 26, 29].

4. Numerical Example

In this section, an example is provided to illustrate the efficiency of this approach. For the sake of comparison purposes, we use the two norm and infinity norm of errors.

Consider the following weakly singular PIDE [21, 28]

$$u_t(x, t) = \int_0^t (t-s)^{-\frac{1}{2}} u_{xx}(x, s) ds, \quad 0 < x < 1, \quad 0 \leq t \leq T,$$

with the boundary and initial conditions

$$u(0, t) = u(1, t) = 0, \quad 0 \leq t \leq T,$$

$$u(x, 0) = \sin(x), \quad 0 \leq x \leq 1.$$

The exact solution is $u(x, t) = M(\pi^2 t^{\frac{5}{2}}) \sin(\pi x)$, where M denotes the function

$$M(z) = \sum_{n=0}^{\infty} (-1)^n \Gamma\left(\frac{3}{2}n + 1\right)^{-1} z^n.$$

We will use $\theta = \frac{1}{2}$, $h = 0.001$, $T = 1$, and $N = 25$.

Errors of the numerical solutions for TPS-RBF ($m = 4$), IMQ-RBF ($c = 0.1$), and Sech-RBF ($c = 0.1$) are shown in the Table 2 and are plotted in Figures 1, 2, and 3, respectively.

Table 2. Errors.

t	TPS-RBF		IMQ-RBF		Sech-RBF	
	$\ E\ _{\infty}$	$\ E\ _2$	$\ E\ _{\infty}$	$\ E\ _2$	$\ E\ _{\infty}$	$\ E\ _2$
0.1	5.98e-04	2.16e-03	8.92e-03	2.71e-02	5.31e-03	1.60e-02
0.2	1.83e-03	6.34e-03	6.46e-03	2.48e-02	3.16e-03	1.25e-02
0.3	2.47e-03	8.53e-03	5.36e-03	1.70e-02	2.29e-03	7.10e-03
0.4	2.17e-03	7.51e-03	3.65e-03	7.75e-03	2.26e-03	5.58e-03
0.5	1.22e-03	4.22e-03	4.06e-03	1.72e-02	2.51e-03	1.18e-02
0.6	1.33e-04	4.46e-04	5.77e-03	2.31e-02	3.52e-03	1.41e-02
0.7	6.64e-04	2.31e-03	5.64e-03	2.09e-02	3.15e-03	1.18e-02
0.8	9.82e-04	3.41e-03	3.78e-03	1.34e-02	1.92e-03	6.83e-03
0.9	8.72e-04	3.02e-03	1.41e-03	4.64e-03	5.35e-04	1.72e-03
1.0	5.19e-04	1.80e-03	4.89e-04	2.17e-03	4.76e-04	1.92e-03

5. Conclusion

Three different RBFs are implemented in a FD method for solving a PIDE with a weakly singular kernel successfully. The results of applying the method on the illustrative example confirm the ability and the usefulness of the proposed approach. In comparison with those results reported in [21, 28], this method achieved more accurate results with less data grid points.




Figure 1. TPS-RBF Error.




Figure 2. IMQ-RBF Error.




Figure 3. Sech-RBF Error.

References

- [1] Z. Avazzadeh, Z. Beygi Rizi, F.M. Maalek Ghaini, and G.B. Loghmani. *A numerical solution of nonlinear parabolic-type volterra partial integro-differential equations using radial basis functions*. Eng. Anal. Bound. Elem., Vol. 36, No. 5, 2012, pp. 881–893.
- [2] N.Yu. Bakaev. *On the galerkin finite element approximations to multi-dimensional differential and integro-differential parabolic equations*. BIT Numer. Math., Vol. 37, No. 2, 1997, pp. 237–255.
- [3] M.D. Buhmann. *Multivariable interpolation using radial basis functions*. PhD thesis, University of Cambridge, 1989.
- [4] M.D. Buhmann. *Radial basis functions: theory and implementations*. Cambridge: Cambridge University Press; 2003.
- [5] C. Chen, V. Thomée, and L.B. Wahlbin. *Finite element approximation of a parabolic integro-differential equation with a weakly singular kernel*. Math. Comput., Vol. 58, No. 198, 1992, pp. 587–602.
- [6] R.M. Christensen and L.B. Freund. *Theory of viscoelasticity*. J. Appl. Mech., Vol. 38, No. 3, 1971, pp. 720.
- [7] M. Dehghan and A. Shokri. *A numerical method for two-dimensional schrodinger equation using collocation and radial basis functions*. Comput. Math. Appl., Vol. 54, No. 1, 2007, pp. 136–146.
- [8] M. Dehghan and A. Shokri. *A numerical method for solution of the two-dimensional sine-gordon equation using the radial basis functions*. Math. Comput. Simul., Vol. 79, No. 3, 2008, pp. 700–715.
- [9] L.M. Delves and J.L. Mohamed. *Computational Methods for Integral Equations*. New York: Cambridge University Press; 1985.
- [10] F. Fakhar-Izadi and M. Dehghan. *Space-time spectral method for a weakly singular parabolic partial integro-differential equation on irregular domains*. Comput. Math. Appl., Vol. 67, No. 10, 2014, pp. 1884–1904.
- [11] C. Franke and R. Schaback. *Solving partial differential equations by collocation using radial basis functions*. Appl. Math. Comput., Vol. 93, 1997, pp. 73–82.
- [12] M.E. Gurtin and A.C. Pipkin. *A general theory of heat conduction with finite wave speeds*. Arch. Ration. Mech. An., Vol. 31, No. 2, 1968, pp. 113–126.
- [13] R.L. Hardy. *Multiquadric equations of topography and other irregular surfaces*. J. Geo-phys. Res., Vol. 76, No. 8, 1971, pp. 1905–1915.
- [14] W.J. Hrusa and M. Renardy. *On a class of quasilinear partial integro differential equations with singular kernels*. J. Differ. Equations, Vol. 4, No. 2, 1986, pp. 195–220.
- [15] I.R.H. Jackson. *Radial basis function methods for multivariable approximation*. PhD thesis, University of Cambridge, 1988.

- [16] E.J. Kansa. *Multiquadrics a scattered data approximation scheme with applications to computational fluid-dynamics II, solutions to parabolic, hyperbolic and elliptic partial differential equations*. Comput. Math. Appl., Vol. 19, No. 89, 1990, pp. 147–161.
- [17] C.H. Kim and U.J. Choi. *Spectral collocation methods for a partial integro-differential equation with a weakly singular kernel*. J. Austral. Math. Soc. Ser. B. Appl. Math., Vol. 39, No. 3, 1998, pp. 408–430.
- [18] E. Larsson and B. Fornberg. *A numerical study of some radial basis function based solution methods for elliptic {PDEs}*. Comput. Math. Appl., Vol. 46, No. 56, 2003, pp. 891–902.
- [19] W. Li and X. Da. *Finite central difference/finite element approximations for parabolic integro-differential equations*. Computing, Vol. 90, No. 3-4, 2010, pp. 89–111.
- [20] W. Long, D. Xu, and X. Zeng. *Quasi wavelet based numerical method for a class of par-tial integro-differential equation*. Appl. Math. Comput., Vol. 218, No. 24, 2012, pp. 11842–11850.
- [21] M. Luo, D. Xu, and L. Li. *A compact difference scheme for a partial integro-differential equation with a weakly singular kernel*. Appl. Math. Model., Vol. 39, No. 2, 2015, pp. 947–954.
- [22] R.K. Miller. *An integro differential equation for rigid heat conductors with memory*. J. Math. Anal. Appl., Vol. 66, No. 2, 1978, pp. 313–332.
- [23] A.S. Vasudeva Murthy and J.G. Verwer. *Solving parabolic integro-differential equations by an explicit integration method*. J. Comput. Appl. Math., Vol. 39, No. 1, 1992, pp. 121–132.
- [24] R.B. Platte and T.A. Driscoll. *Computing eigenmodes of elliptic operators using radial basis functions*. Comput. Math. Appl, 2004.
- [25] M. Renardy. *Mathematical analysis of viscoelastic flows*, volume 73. Siam, 2000.
- [26] R. Schaback. *Approximation by radial basis functions with finitely many centers*. Constr. Approx., Vol. 12, No. 3, 1996, pp. 331–340.
- [27] F. Shakeri and M. Dehghan. *A high order finite volume element method for solving elliptic partial integro-differential equations*. Appl. Numer. Math., Vol. 65, 2013, pp. 105–118.
- [28] T. Tang. *A finite difference scheme for partial integro-differential equations with a weakly singular kernel*. Appl. Numer. Math., Vol. 11, No. 4, 1993, pp. 309–319.
- [29] H. Wendland. *Error estimates for interpolation by compactly supported radial basis functions of minimal degree*. J. Approx. Theory, Vol. 93, No. 2, 1998, pp. 258–272.
- [30] E.G. Yanik and G. Fairweather. *Finite element methods for parabolic and hyperbolic partial integro-differential equations*. Nonlinear Anal–Theor, Vol. 12, No. 8, 1998, pp. 785–809.
- [31] H. Zhang, X. Han, and X. Yang. *Quintic B-spline collocation method for fourth order partial integro-differential equations with a weakly singular kernel*, Appl. Math. Comput., Vol. 219, 2013, pp. 6565–6575.
- [32] X. Yang, D. Xu, and H. Zhang, *Crank–Nicolson/quasi-wavelets method for solving fourth order partial integro-differential equation with a weakly singular kernel*, J. Comput. Phys., Vol. 234, 2013, pp. 317–329.
- [33] M. Q. Wang, C. X. Wang, Ming Li and C. S. Chen, *A Numerical Scheme Based on FD-RBF to Solve Fractional-Diffusion Inverse Heat Conduction Problems*, Numer. Heat TR A-Appl., Vol 68, No, 9, 2015, pp. 978-992.
- [34] M.K. Kadalbajoo, A. Kumar, and L.P. Tripathi *A radial basis functions based finite differences method for wave equation with an integral condition*, Appl. Math. Comput. Vol. 253, 2015, pp. 8–16.